

SENSISpec ELISA Pistachio 96/48 Tests

Enzyme Immunoassay for the Quantitative Determination of Pistachio in Food (Cat.nr. HU0030021, HU0030045)

Sensitivity (Pistachio)	0.00-0.84 ppm
Recovery	70-96%
Incubation Time	60 min

1. GENERAL INFORMATION

Pistachio (*Pistacia vera*) belongs to the family of Anacardiaceae. With about 21% the fraction of proteins in pistachio seed is very high. Some of these proteins, like the 2S albumin Pis v 1, the 11S globulin Pis v 2 or the 7S vicilin Pis v 3 are known for being allergenic. Many of them are heat resistant making them stable to different production processes. For this reason, pistachio represents an important food allergen.

For pistachio-allergic persons, hidden pistachio allergens in food are a critical problem. Already very low amounts of pistachio can cause allergic reactions, which may lead to anaphylactic shock in severe cases. Because of this, pistachio-allergic persons must strictly avoid the consumption of pistachio-containing food. Cross-contamination, mostly in consequence of the production process, is often noticed. This explains why in many cases the existence of pistachio residues in food cannot be excluded. For this reason, sensitive detection systems for pistachio residues in foodstuffs are required.

The **Eurofins SENSISpec Pistachio ELISA** represents a highly sensitive detection system and is particularly capable of the quantification of pistachio residues in cookies, ice cream, sweets, chocolate, spices and food supplements. Furthermore, it is validated for rinse water / CIP s and swab samples.

2. PRINCIPLE OF THE TEST

The **Eurofins SENSISpec Pistachio** quantitative test is based on the principle of the enzyme linked immunosorbent assay. An antibody directed against pistachio proteins is bound on the surface of a microtiter plate. Pistachio containing samples or standards are given into the wells of the microtiter plate. After 20 minutes incubation at room temperature, the wells are washed with a diluted washing solution to remove unbound material. A peroxidase conjugated secondary antibody directed against pistachio proteins is given into the wells and after 20 minutes of incubation the plate is washed again. A substrate solution is added and incubated for 20 minutes, resulting in the development of a blue color. The color development is inhibited by the addition of a stop solution, and the color turns yellow.

The yellow color is measured photometrically at 450 nm. The concentration of pistachio is directly proportional to the color intensity of the test sample.

3. PRECAUTIONS

Full compliance of the following good laboratory practices (GLP) will determine the reliability of the results:

- 1) Prior to beginning the assay procedure, bring all reagents to room temperature (20-25°C).
- 2) All reagents should be mixed by gentle inversion or swirling prior to use. Do not induce foaming.
- 3) Once the assay has been started, all subsequent steps should be completed without interruption and within the recommended time limits.
- 4) Replace caps in all the reagents immediately after use. Do not interchange vial stoppers.
- 5) Use a separate disposable tip for each specimen to prevent cross-contamination.
- 6) All specimens and standards should be run at the same time, so that all conditions of testing are the same.
- 7) Do not mix components from different batches.
- 8) Do not use reagents after the expiration date.
- 9) Check both precision and accuracy of the laboratory equipment used during the procedure (micropipette, ELISA reader etc.).

4. HEALTH AND SAFETY INSTRUCTIONS

- 1) Do not smoke or eat or drink or pipet by mouth in the laboratory.
- 2) Wear disposable gloves whenever handling patient specimens.
- 3) Avoid contact of substrate and stop solution with skin and mucosa (possible irritation, burn or toxicity hazard). In case of contact, rinse the affected zone with plenty of water.
- 4) Handling and disposal of chemical products must be done according to good laboratory practices (GLP).

5. REAGENTS

The kit contains reagents for 96/48 determinations. They have to be stored at 2-8°C. Expiry dates are found on the labels of the bottles and the outer package.

- 1) Microtiter plate consisting of 12/6 strips with 8 breakable wells each, coated with anti-pistachio antibodies.
- 2) Pistachio Standards (0; 1; 4; 10; 40 ppm of pistachio): 5 vials with 2.0 mL each, dyed red, ready-to-use
- 3) Conjugate (anti-pistachio-peroxidase): 15/7.5 mL, dyed red, ready-to-use.
- 4) Substrate Solution (TMB): 15 mL, ready-to-use.
- 5) Stop Solution (0.5 M H₂SO₄): 15 mL, ready-to-use.
- 6) Extraction and sample dilution buffer (Tris): 2/1 x 120 mL as 10x concentrate, dyed red. Dilute 1+9 with distilled water. Stored at 4°C the diluted buffer is stable for at least one week. If during the cold storage crystals precipitate, the concentrate should be warmed up to 37°C for 15 minutes.
- 7) Washing Solution (PBS + Tween 20): 60 mL as 10x concentrate. Dilute 1+9 with distilled water. Stored at 4°C the diluted buffer is stable for at least 4 weeks. If during the cold storage crystals precipitate, the concentrate should be warmed up to 37°C for 15 minutes.
- 8) Plastic bag to store unused microtiter strips.
- 9) Instruction Manual.

6. ADDITIONAL INSTRUMENTATION AND REAGENTS (NOT PROVIDED)

6.1 Instrumentation

- 100 - 1000 µL micropipettes
- Volumetric flask
- Analytical balance
- Mortar, mixer
- Water bath
- Centrifuge
- ELISA reader (450 nm)

6.2 Reagents

- double distilled water
- Polyvinylpyrrolidone (PVP), optional

7. SAMPLE PREPARATION

Due to the high risk of cross-contamination, all applied instruments like applicator, mortar, glass vials etc. have to be **cleaned thoroughly** before and after each sample. Pistachio proteins adhere very strongly to different surfaces. In certain cases, they can resist common dishwasher cleaning.

To identify possible cross-contamination caused by previous extractions it is strongly recommended to note the sequence of the extractions.

The following sample preparation should be applied for all solid food samples:

- 1) To maximize homogeneity and representativeness of the sample drawing, a minimum of 5 g sample should be pulverized finely in a mortar, impact mill, etc.
- 2) 1 g of the homogenized mixture is suspended in 20 mL of **pre-diluted** extraction and sample dilution buffer. Afterwards the suspension is incubated for 15 min in a preheated water bath at 60°C. To ensure good homogeneity, the samples should be shaken every two minutes.
- 3) The samples are centrifuged for 10 minutes at 2000 g. If it is not possible to separate the supernatant from the precipitate completely, the suspension should be filtrated if necessary.
- 4) 100 µL of particle-free solution are applied per well. If the results of a sample are out of the measuring range, further dilution with the **pre-diluted** extraction and sample dilution buffer is necessary. The additional dilution has to be considered when calculating the concentration.

The following sample preparation should be applied for liquid food samples:

1 mL of liquid sample is diluted in 19 mL of **pre-diluted** extraction and sample dilution buffer. Afterwards the suspension is incubated for 15 min in a preheated water bath at 60°C. To ensure good homogeneity, the samples should be shaken every two minutes. The process is continued at point 3 of the solid sample extraction process.

The following variation should be applied for polyphenol containing food samples like chocolate:

Dilute 1 g of Polyvinylpyrrolidone (PVP) in 100 mL of **pre-diluted** extraction buffer. Apply the buffer as extraction buffer in the sample preparations stated above


The following sample preparation should be applied for rinse water samples:

- 1) Adjust the pH of the sample to 8.2 (+/- 0.5)
- 2) 1 mL of liquid sample is diluted in 4 mL of **pre-diluted** extraction and sample dilution buffer. The process is continued at point 4 of the solid food sample extraction process.

The following sample preparation should be applied for swab samples on dry surfaces:

- 1) Mark out 5x5 cm area or use swab directly on (e.g. uneven) area.

- 2) Moisten the swab in 1 mL **pre-diluted** extraction and sample dilution buffer previously applied in a test tube.
- 3) Swab marked area by using crosshatch (1. horizontally, 2. vertically, 3. diagonally) technique while rotating the tip.



Place the swab into the test tube.

- 4) Shake the test tube for 1 minute to release the sample from the swab. The process is continued at point 4 of the solid food sample extraction process.

For wet surfaces exactly the same procedure is applied without prior need to moisten the swab.

8. PROCEDURE

The washing solution is supplied as 10x concentrate and has to be **diluted** 1+9 with double distilled water before use.

In any case the **ready-to-use** standards provided should be determined twofold. When samples in great quantities are determined, the standards should be pipetted once before the samples and once after the samples. For final interpretation the arithmetic mean is used for calculation.

In consideration of GLP and quality control requirements, a duplicate measurement of samples is recommended.

The procedure is according to the following scheme:

- 1) Prepare samples as described above.
- 2) Pipet 100 µL ready-to-use standards or prepared samples in duplicate into the appropriate wells of the microtiter plate.
- 3) Incubate for 20 minutes at room temperature.
- 4) Wash the plate three times as follows: Discard the contents of the wells (dump or aspirate). Pipet 300 µL of diluted washing solution into each well. After the third repetition, empty the wells again and remove residual liquid by striking the plate against a paper towel. The wash procedure is critical. Insufficient washing will result in poor precision and falsely elevated absorbencies.
- 5) Pipet 100 µL of conjugate (anti-pistachio-peroxidase) into each well.
- 6) Incubate for 20 minutes at room temperature.
- 7) Wash the plate as outlined in 4.
- 8) Pipet 100 µL of substrate solution into each well.

- 9) Allow the reaction to develop in the dark (e.g. cupboard or drawer; the chromogen is light-sensitive) for 20 minutes at room temperature.
- 10) Stop enzyme reaction by adding 100 µL of stop solution (0.5 M H₂SO₄) into each well. The blue color will turn yellow upon addition.
- 11) After thorough mixing, measure absorbance at 450 nm (reference wavelength 620 nm), using an ELISA reader. The color is stable for 30 minutes.

9. CALCULATION OF RESULTS

The following evaluation procedure should be applied for all **food samples** prepared by the procedure as stated *Sample Preparation*:

The ready-to-use standards are prepared for a direct determination of food sample concentrations. The dilution (1:20) of samples in the extraction process as described in the above stated sample preparation procedure is already considered. Additional dilution due to high sample concentration has to be accounted for.

- 1) Calculate the average optical density (OD 450 nm) for each set of reference standards or samples.
- 2) Construct a standard curve by plotting the mean optical density obtained for each reference standard against its concentration in ppm on semi-log graph paper with the optical density on the vertical (y) axis and the concentration on the horizontal (x) axis. Alternatively, the evaluation can be carried out by software. In this case the 4-parameter method should be preferred.
- 3) Using the mean optical density value for each sample, determine the corresponding concentration of pistachio in ppm from the standard curve. Depending on experience and/or the availability of computer capability, other methods of data reduction may be employed.

The following evaluation procedure should be applied for **rinse water samples** prepared according to the procedure stated *Sample Preparation*:

- 1) Apply the evaluation procedure for food samples as stated above.
- 2) Divide the result by 4 in order to compensate for the different dilution factor of the extraction procedure to receive the sample concentration in mg/L.

The following evaluation procedure should be applied for **swab samples** prepared according to the procedure stated in *Sample Preparation*:

- 1) Apply the evaluation procedure for food samples as stated above.

- 2) Multiply the result (ppm) by 2 in order to compensate for the different dilution factor of the extraction procedure to receive the sample concentration in ng/cm².

10. TYPICAL STANDARD VALUES

The following table contains an example for a typical standard curve. The binding is calculated as percent of the absorption of the 40 ppm standard. These values are only an example and should not be used instead of the standard curve which has to be measured in every new test.

Pistachio (ppm)	% binding of 40 ppm
40	100
10	65
4	39
1	15
0	3

11. PERFORMANCE

11.1 Sensitivity

The limit of detection (LOD) of the **Eurofins SENSISpec Pistachio test** is 0.07ppm pistachio.

Validation experiments with common matrices resulted in the following mean LODs [ppm].

Cookies	0.18
Ice-cream	0.06
Sweets	0.02
Chocolate	0.15
Spices	0.46
Food Supplement	0.13

The limit of quantification (LOQ) of the **Eurofins SENSISpec Pistachio test** is 1 ppm pistachio.

Due to the variety of sample matrices and their influence on the blank, results less than the LOQ should be treated as negative.

11.2 Precision

Intra-Assay Precision	6.4%
Inter-Assay Precision	7.8%
Inter-Extraction Precision	5.8%

1.3 Linearity

The serial dilution of spiked samples (cookies, cereals, ice cream, chocolate, sweets and food supplements) resulted in a mean dilution linearity of 97-119%.

11.4 Cross-reactivity

For the following foods no cross-reactivity could be detected:

Adzuki bean	Cow' milk	Paprika
Almond	Cumin	Pea
Apricot	Dill	Peach
Barley	Duck	Peanut
Bean, white	Egg, dried	Pecan
Beef	Fennel	Pepper, black
Bovine gelatin	Fenugreek	Pine seed
Brazil nut	Garden cress	Pork
Buckwheat	Garlic, fresh	Potato
Cabbage, white	Garlic, granul.	Prawn
Caraway	Ginger, fresh	Pumpkin seed
Cardamom	Ginger, ground	Radish
Carob gum	Gliadin	Rice
Carrot	Goat's milk	Rye
Cayenne	Guar gum	Saccharose
Celery	Gum arabic	Shrimps
Cherry	Hazelnut	Soy flour
Chestnut	Kidney bean	Soy lecithin
Chia	Kiwi	Split pea
Chicken	Lamb	Sunflower seed
Chickpea	Leek	Thyme
Chili	Lentil	Tofu
Cinnamon	Lupin	Tomato
Clove	Macadamia	Turkey
Cocoa	Nutmeg	Turmeric
Coconut	Oats	Walnut
Cod	Onion	Wheat
Corn		

The following cross reactions were determined:

Poppy seed	0.0001%
Rapeseed	0.0002%
Sesame	0.0001%
Horseradish	0.0001%
Flaxseed	0.0001%
Cashew	0.0033%
Mustard, yellow	0.0002%

11.5 Recovery

Mean recovery was determined by spiking samples with different amounts of pistachio:

Cookies	90%
Sweets	78%
Ice-cream	86%
Chocolate	80%
Spices	80%
Food Supplement	72%

12. LIABILITY

Eurofins Technologies Hungary shall not be liable for any damages to the customer caused by the improper use of the kit and for any action undertaken as a consequence of the results.

Eurofins Technologies Hungary shall not be liable for the unsafe use of the kit out of the current European safety regulations.